

ORIGINAL INVESTIGATION

Open Access



DPP-4 inhibition has no acute effect on BNP and its N-terminal pro-hormone measured by commercial immune-assays. A randomized cross-over trial in patients with type 2 diabetes

Gian Paolo Fadini*, Benedetta Maria Bonora, Mattia Albiero, Martina Zaninotto, Mario Plebani and Angelo Avogaro

Abstract

Background: Use of dipeptidyl peptidase-4 inhibitors (DPP4-i) for the treatment of type 2 diabetes (T2D) has been associated with a possible increase in the risk for heart failure (HF). B-type natriuretic peptide (BNP), which is both a biomarker of HF and a hemodynamically active hormone, is a substrate of DPP-4. We herein tested the acute effects of the DPP-4i linagliptin on BNP and NT-proBNP in a cross-over placebo-controlled trial in patients with T2D with and without chronic kidney disease (CKD).

Methods: B-type natriuretic peptide and NT-proBNP were measured using commercially available clinical-grade immune-assays at baseline and at the end of a 4-day treatment with placebo and linagliptin. Changes from baseline during each treatment arm, as well as placebo-subtracted effects of linagliptin on BNP and NT-proBNP were calculated.

Results: 46 patients completed the study, 18 of whom were affected by CKD. Baseline BNP and NT-proBNP levels increased with age, were elevated in CKD patients, and inversely correlated with estimated glomerular filtration rate. No significant change was detected in BNP and NT-proBNP levels after treatment with linagliptin or placebo in patients with or without CKD. Only in CKD patients the placebo-subtracted effect of linagliptin indicated a significant reduction in NT-proBNP levels, but this finding was not statistically robust.

Conclusions: Acute treatment with a DPP-4i exerts no clinically-meaningful effects on BNP and NT-proBNP. As routinely used immunoassays do not discriminate between intact/active and cleaved BNP, these data cannot rule out an effect of DPP-4i on HF pathophysiology.

Trial registration NCT01617824

Keywords: Heart failure, Enzyme, Proteases, Linagliptin, Kidney disease

Background

Dipeptidyl peptidase-4 inhibitors (DPP-4i) are routinely used for the treatment of type 2 diabetes (T2D). Their

glucose-lowering activity results from inhibition of the enzymatic cleavage of glucagon-like peptide (GLP)-1 and glucose insulinotropic peptide (GIP) by DPP-4 [1]. DPP-4 has several other physiological substrates, including cytokines, chemokines, and neuro-hormones, providing the biological basis for non-glycemic effects of DPP-4i [2]. However, comparatively fewer peptides have been

*Correspondence: gianpaolofadini@hotmail.com; gianpaolo.fadini@unipd.it
Department of Medicine, University of Padova, Via Giustiniani,2,
35128 Padua, Italy

identified as endogenous physiological substrates in vivo [3].

Concerns have been raised on the possibility that DPP-4i increase the risk of heart failure (HF), but the mechanisms are largely unknown [4, 5]. Intact B-type natriuretic peptide (BNP₁₋₃₂) is cleaved by DPP-4, generating BNP₃₋₃₂ [6]. BNP, which is produced by cardiomyocytes in response to hemodynamic stress and neuro-hormonal stimulation, is a clinical-grade biomarker of HF [7]. In turn, BNP is involved in the pathophysiology of HF, as it induces vasodilation and natriuresis, thereby antagonizing the effects of angiotensin-II [8, 9]. BNP concentrations are reduced in people with obesity, insulin resistance, and diabetes, and this deficiency may contribute to their cardiovascular risk [10]. Thus DPP-4i may exert beneficial effects on cardiac function, by increasing the proportion of intact/active BNP₁₋₃₂ versus cleaved BNP₃₋₃₂, in addition to GLP-1 mediated cardioprotection [11].

BNP₁₋₃₂ derives from proBNP (108 amino-acid) after removal of an N-terminal (NT) fragment of 76 amino-acids by pro-hormone convertases [12]. Although BNP and NT-proBNP are released at equimolar concentrations, the half-life of the NT-proBNP in the circulation is longer, resulting in higher concentrations. In patients with diabetes, despite a possible reduction of BNP, the clinical predictive capacity of NT-proBNP has been shown to be preserved [13, 14]. Interestingly, also proBNP₁₋₁₀₈ and NT-proBNP₁₋₇₆ are candidate substrates of the enzymatic activity of DPP-4, as they have proline in the second N-terminal position, where the exopeptidase activity of DPP-4 locates. Commercially available immuno-assays are presumably unable to distinguish

between BNP₁₋₃₂ and BNP₃₋₃₂ [15], and they may even detect proBNP_{1-108/3-108}, but not NT-proBNP [16]. As compared to BNP₁₋₃₂, truncated BNP₃₋₃₂ appears to have equal cGMP activating properties in vitro [17], but lower activity in vivo [18], probably because of a higher susceptibility to degradation by other peptidases. Specificity of NT-proBNP immunoassays is unknown, but epitopes recognized by monoclonal antibodies do not appear to span the first 2 N-terminal residues. Based on these considerations, the net effect of a DPP-4i therapy on diagnostic BNP and NT-proBNP determinations is unpredictable (Fig. 1).

Available data on the effects of DPP-4i on proBNP-derived peptides in T2D mostly come from large clinical trials wherein NT-proBNP levels were measured years after therapy with a DPP-4i or placebo [19, 20]. Rather, if DPP-4i has direct effects on proBNP processing, this should be detectable within a short time frame. To address the acute effects of DPP-4i on BNP and NT-proBNP, we used samples from a placebo-controlled cross-over trial testing the effects of a 4-day therapy with the DPP-4i inhibitor linagliptin on humoral factors [21]. Differently from other DPP-4i, linagliptin has no renal excretion, and is thereby particularly suitable for the treatment of patients with chronic kidney disease (CKD) [22, 23], which is a major risk factor for HF [24].

Methods

Study design

The NCT01617824 was a randomized, single-blind, placebo-controlled, cross-over study designed to test the acute effects of the DPP-4 inhibitor linagliptin on cytokines, hormones, and inflammatory mediators. The




Fig. 1 Sequential cleavage of proBNP to originate BNP, NT-proBNP and their by-products. Biological activity on cardiac function is reported. DPP-4 can cleave the 2 N-terminal residues of proBNP, BNP, and NT-proBNP, generating inactive or less active peptides

primary study results have been published before [21]. Briefly, T2D patients with or without CKD, received a 4-day treatment with linagliptin 5 mg and placebo in a random order with a 14-day wash-out period. Before and at the 5th day of each treatment period, fasting blood samples were drawn. Aliquots of EDTA and heparin plasma were separated and stored at -80°C until analyses. CKD was defined as an estimated glomerular filtration rate (eGFR) of less than 60 ml/min/1.73 m², based on the CKD-EPI formula [25] and graded according to the Kidney Disease Outcomes Quality Initiative (KDOQI) [26]. With this design, we were able to detect significant changes in DPP-4 activity, levels of intact (uncleaved) GLP-1 and SDF-1 α , along with an increase in CD34⁺KDR⁺ cells, reflecting a biological consequence of elevated SDF-1 α [21], an effect that was observed also for saxagliptin [27]. For this study, untouched frozen aliquots of plasma were thawed and used for the quantification of BNP and NT-proBNP.

Analytical methods

B-type natriuretic peptide was quantified in EDTA plasma using a chemiluminescent, microparticle-capture immunoassay (Abbott Diagnostics kit, #8K28) on the modular automated ARCHITECT iSystem platform. The range of plasma BNP concentrations revealed using this assay is 10–5000 pg/ml. The coefficient of variation (CV), as reported by the manufacturer, is <5%. This BNP assay results in no cross-reactivity with ANP, Angiotensin-I, Angiotensin-II, Angiotensin-III, CNP, and NT-proBNP.

NT-proBNP was quantified in heparinised plasma using a solid-phase two-site chemiluminescent immunometric assay (Siemens IMMULITE 1000 Turbo). The sensitivity of this assay is 15 pg/ml, with a reportable range up to 35,000 pg/ml. The CV reported by the manufacturer is 9%. No cross-reactivity has been detected with ANP, NT-proANP, BNP, CNP, Adrenomedullin, Angiotensin-I, Angiotensin-II, Angiotensin-III, Endothelin, Renin, Urodilatin, and Arg-Vasopressin.

Statistical analysis

Data are expressed as mean \pm standard error if normal or as median (interquartile range) if not normal. Normality was checked using the Shapiro–Wilk test and non-normal variables were log-transformed before analysis. Within-group changes in continuous variables were analyzed using the paired two-tail Student's *t* test. For each patients in each group of treatment order, we calculated the effect of placebo, the effect of linagliptin, and the placebo-subtracted effect of linagliptin. The generalized linear model (GLM) was used to analyze the effect of treatment and order by the cross-over design. Statistical

significance was accepted at $p < 0.05$ and SPSS version 22.0 was used.

Sample size was originally chosen to achieve a 80% power to detect a significant difference in the primary end-point (a difference in circulating CD34⁺KDR⁺ cells), which was fully satisfied. Based on within-patients standard deviations of 37% for BNP (26 pg/ml) and 40% for NT-proBNP (227 pg/ml), we calculated a priori that this study had 80% power to detect a treatment difference at a two-sided 0.05 significance level, if the true difference between treatments was 22% for BNP (15 pg/ml) and 24% for NT-proBNP (136 pg/ml).

Results

Characteristics of study patients

A total of 46 patients completed the study. Detailed baseline clinical characteristics of the participants have been reported previously [21] and are herein summarized in Table 1. There was no difference between patients randomized to the placebo-linagliptin ($n = 22$) or the linagliptin-placebo ($n = 24$) treatment order. No mild or severe adverse event was reported during treatment or wash-out and no change in fasting metabolic variables (glucose, triglycerides and fatty acids) was observed [21].

Baseline values of BNP and NT-proBNP

As the distributions of BNP and NT-proBNP were highly skewed, data are presented as median (IQR) and values were log-transformed before statistical testing.

The median baseline plasma BNP level was 20.4 pg/ml (IQR 10.0–43.3). BNP was below threshold (10 pg/ml) in $n = 15$ patients (32.6%) and was above the decisional cut-off (100 pg/ml) [28] in 6 patients (13.0%). BNP levels increased with age ($r = 0.40$; $p = 0.003$), were higher in patients with CKD than in those without (43.1 [IQR 22.1–98.5] versus 12.5 [IQR 10.0–23.0] pg/ml; $p = 0.0022$) and were inversely correlated with eGFR ($r = -0.45$; $p < 0.001$).

The median baseline NT-proBNP level was 101.0 pg/ml (IQR 35.3–314.8) and $n = 5$ patients (10.9% had values above the decisional cut-off (900 pg/ml) [28]. NT-proBNP levels increased with age ($r = 0.52$; $p < 0.001$), were higher in patients with CKD than in those without (238.5 [IQR 115.0–554.8] versus 44.0 [IQR 24.3–101.0] pg/ml; $p < 0.001$) and were inversely correlated with eGFR ($r = -0.50$; $p < 0.001$).

Levels of BNP and NT-proBNP were highly correlated ($r = 0.94$).

Effects of DPP-4 inhibition on BNP and NT-proBNP

Overall, no significant change versus baseline was observed in BNP and NT-proBNP levels after treatment with linagliptin or placebo (Fig. 2). The

Table 1 Baseline characteristics of study patients

Variable	All patients
Number	46
Age, years	63.7 ± 1.3
Sex male, %	71.7
Body mass index, kg/m ²	31.1 ± 0.7
Waist, cm	105.4 ± 2.2
HbA1c, % (mmol/mol)	7.6 ± 0.2 (60 ± 2)
Risk factors	
Smoking habit, %	13.0
Hypertension, %	89.1
Total cholesterol, mg/dl	165.3 ± 5.5
HDL cholesterol, mg/dl	49.9 ± 2.2
LDL cholesterol, mg/dl	91.6 ± 5.1
Triglycerides, mg/dl	119.2 ± 8.5
Albumin/creatinine ratio (mg/g)	129.5 ± 44.1
Creatinine, mg/dl	1.11 ± 0.06
eGFR, ml/min/1.73 m ²	75.5 ± 3.9
Complications	
Retinopathy, %	28.2
Neuropathy, %	17.9
Coronary artery disease, %	30.4
Peripheral arterial disease, %	21.7
Cerebrovascular disease, %	47.8
Medications	
Metformin, %	65.2
Sulphonylurea, %	6.5
Repaglinide, %	4.3
Pioglitazone, %	6.5
Insulin, %	43.4
ACE inhibitors/ARBs, %	76.1
Other anti-hypertensives, %	78.2
Statin, %	80.4
Anti-platelet agents, %	56.5

Data are presented as mean ± standard error, or as percentage, where appropriate. More details can be found in [21]

placebo-subtracted effect of linagliptin on BNP and NT-proBNP were 0.0 pg/ml (IRQ −19.0 to 1.7) and −19.5 pg/ml (IQR −62.3 to 19.3), respectively. In patients with CKD (n = 18) the change of NT-proBNP during linagliptin treatment was not significant (−17.0; IRQ −59.3 to 23.0 pg/ml), but was significantly different from the change during placebo (4.5; IQR −57.8 to 109.3 pg/ml; p = 0.022). This resulted in a significant placebo-subtracted reduction in NT-proBNP attributable to linagliptin treatment (Table 2; Fig. 2). The post hoc power of the study to detect this finding was calculated to be <20%. Two patients with CKD had a previous history of HF, but placebo-subtracted changes of BNP (−72.7 and +34.0 pg/ml) or NT-proBNP (−42 and +216 pg/ml) were inconsistent with an effect of DPP-4i.

No carry-over effect was noted for both BNP and NT-proBNP. No correlation was detected between BNP or NT-proBNP and DPP-4 activity, nor between change in BNP or NT-proBNP and change in DPP-4 activity.

Discussion

We show that therapy with a DPP-4i has no acute effects on BNP and NT-proBNP levels measured with routine diagnostic immuno-assays. This study was not designed to test the acute effects of DPP-4i on cardiac function, but our findings re-assure on the safety of DPP-4i concerning diagnosis and prognostic evaluation of HF.

The clinical relevance of the interplay between DPP-4i and BNP/NT-proBNP levels has emerged after publication of the results of SAVOR-TIMI trial, wherein patients treated with the DPP-4i saxagliptin exhibited a significant 27% excess risk of hospitalization for HF compared to placebo [19, 29]. Meta-analyses of randomized controlled trials were unable to rule out the concern that DPP-4i therapy may favour HF [4, 5]. Real-world data did not confirm an association between DPP-4i and hospitalization for HF [30, 31], nor show adverse prognosis in HF patients treated with DPP-4i [32, 33]. Furthermore, the eventual mechanisms remain elusive. In the SAVOR-TIMI trial, the risk of HF associated with saxagliptin therapy was almost exclusively observed in patients with a baseline NT-proBNP level within the most elevated quartile [19]. During a follow-up of about 2 years, NT-proBNP levels increased in both the placebo and saxagliptin group, but the increase was slightly blunted by saxagliptin [19]. The relevance of this finding is limited because exceeding HF cases in saxagliptin-treated patients were observed only in the first 6 months of therapy [19]. In another placebo-controlled trial conducted on T2D patients after an acute coronary event, the DPP-4i alogliptin was associated with a non-significant increase in the risk of hospitalization for HF [34]. During an average 1.5 year follow-up, NT-proBNP concentrations decreased significantly and similarly in the two groups [20]. Our study shows for the first time that BNP and NT-proBNP are not directly affected by DPP-4i because no acute effect was detected. This suggests that the observed changes in NT-proBNP over the long run most likely reflect the natural history of HF, rather than effects of therapy.

Chronic kidney disease is a one of the strongest risk factors for HF [24]. As BNP and NT-proBNP levels were higher in CKD patients, the amplitude of their excursions after DPP-4i or placebo was also larger. Although consistent with a study showing that linagliptin decreased BNP in an experimental model of uremic cardiomyopathy [35], the modest placebo-subtracted effect of DPP-4i on NT-proBNP reduction we observe in CKD patients




Table 2 BNP and NT-proBNP levels, expressed as median (IQR) during treatment with placebo or linagliptin

	Placebo			Linagliptin			Placebo-subtracted change
	Pre	Post	Change	Pre	Post	Change	
BNP							
All	22.9 (10.0–42.8)	21.3 (10.0–52.6)	0.0 (–0.9 to 15.2)	19.6 (10.0–44.3)	22.6 (10.0–39.3)	0.0 (–2.7 to 7.8)	0.0 (–19.0 to 1.7)
No CKD	11.4 (10.0–23.3)	11.9 (10.0–29.5)	0.0 (–0.9 to 0.3)	10.6 (10.0–21.8)	12.5 (10.0–27.4)	0.0 (0.0–4.6)	0.0 (–1.0 to 1.2)
CKD	38.5 (26.8–80.3)	56.3 (24.6–107.1)	7.3 (–0.7 to 26.0)	44.2 (20.3–129.0)	37.4 (25.0–78.9)	0.0 (–9.5 to 12.1)	–17.6 (–46.8 to 0.8)
NT-proBNP							
All	101.0 (36.5–314.5)	122.0 (38.0–267.0)	2.0 (–32.0 to 30.5)	101.0 (44.0–299.0)	78.0 (36.0–272.0)	–3.5 (–27.5 to 19.3)	–19.5 (–62.3 to 19.3)
No CKD	46.0 (26.0–101.0)	57.0 (20.0–167.0)	0.0 (–12.0 to 20.0)	51.0 (28.5–121.5)	41.0 (26.5–99.0)	3.0 (–17.5 to 17.3)	–2.5 (–35.0 to 15.8)
CKD	218.5 (108.8–554.8)	261.0 (130.0–696.5)	4.5 (–57.8 to 109.3)	238.5 (107.5–611.5)	184.0 (109.0–483.0)	–17.0* (–59.3 to 23.0)	–50.0# (–101.5 to 18.0)

* Significantly different from placebo treatment (p < 0.05 at paired t test on log-transformed data or Mann–Whitney test)

Significantly different from zero

had very low statistical power, was no longer significant after adjusting for multiple testing, and is unlikely to be of any clinical meaning, as NT-proBNP is not biologically active.

The present study has limitations inherent to the small sample size, the fact that a minority of patients had CKD, thereby lowering power in this subgroup, and the lack of data in patients with decompensated HF. Finally, more subtle changes in BNP and NT-proBNP induced by DPP-4i may have been missed, since the study was powered for a minimal detectable change of 22 and 24%, respectively. Since clinical-grade commercially available immuno-assays do not distinguish the intact and cleaved forms of BNP and NT-proBNP, our data provide no clear indication of whether DPP-4i interferes with the *in vivo* processing of the two peptides, and whether it intervenes in the pathophysiology of HF. However, any eventual significant change in the relative proportion of BNP_{1–32} and BNP_{3–32} or in the proportion of NT-proBNP_{1–76} and NT-proBNP_{3–76} induced by DPP-4i may nonetheless result in modifications of immune-reactive (total) BNP and NT-proBNP levels, respectively. This has been shown for GLP-1 and SDF-1 α [21], possibly reflecting compensatory secretion and/or changes in sequential cleavage by different peptidases. Furthermore, experimental studies suggest that linagliptin exerts favourable effects on ischemia–reperfusion injury [36], which in the long-term can translate into protection from HF.

Conclusion

Although exact discrimination of the various proBNP-derived peptides will require sophisticated, time consuming and costly mass spectrometric approaches [15], data obtained with diagnostic assays indicate that DPP-4i has no clinically appreciable effects on BNP and NT-proBNP. Further studies will be needed to dissect whether DPP-4i interferes with the biological action of BNP and whether this is linked to HF risk in patients with T2D.

Abbreviations

ANP: atrial natriuretic peptide; BMI: body mass index; BNP: B-type natriuretic peptide; CKD: chronic kidney disease; CNP: cardiac natriuretic peptide; DPP-4: dipeptidyl peptidase-4; EDTA: ethylenediamineacetic acid; eGFR: estimated glomerular filtration rate; GIP: glucose-insulinotropic peptide; GLM: generalized linear model; GLP-1: glucagon-like peptide 1; HF: heart failure; IQR: interquartile range; KDOQI: kidney disease outcomes quality initiative; NT: N-terminal; SDF: stromal derived factor; T2D: type 2 diabetes.

Authors' contributions

GPF study design and conduction, data collection, analysis and interpretation, manuscript writing. BMB, study conduction, data collection and analysis, manuscript revision. MA, MZ, MP data collection and analysis, manuscript revision. AA study design, data analysis and interpretation, manuscript writing. All authors read and approved the final manuscript.

Acknowledgements

None.

Competing interests

GPF and AA report receiving lecture fees and honoraria from manufacturer of Linagliptin and other DPP-4 inhibitors. The other authors report no competing interests.

Availability of data and materials

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The study was approved by the Ethical Committee of the University Hospital of Padova (prot. number). All participants provided written informed consent.

Funding

The study was supported by Boehringer-Ingelheim, which also provided the investigational medicinal product. The sponsor had no role in study design, data collection, analysis and interpretation.

Received: 4 January 2017 Accepted: 3 February 2017

Published online: 10 February 2017

References

- Drucker DJ, Nauck MA. The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *Lancet*. 2006;368(9548):1696–705.
- Fadini GP, Avogaro A. Cardiovascular effects of DPP-4 inhibition: beyond GLP-1. *Vasc Pharmacol*. 2011;55(1–3):10–6.
- Vanderheyden M, Bartunek J, Goethals M, Verstreken S, Lambeir AM, De Meester I, Scharpe S. Dipeptidyl-peptidase IV and B-type natriuretic peptide. From bench to bedside. *Clin Chem Lab Med*. 2009;47(3):248–52.
- Elgendy IY, Mahmoud AN, Barakat AF, Elgendy AY, Saad M, Abuzaid A, Wayangankar SA, Bavry AA. Cardiovascular Safety of Dipeptidyl-Peptidase IV Inhibitors: a meta-analysis of placebo-controlled randomized trials. *Am J Cardiovasc Drugs*. 2016. doi:10.1007/s40256-016-0208-x.
- Li L, Li S, Deng K, Liu J, Vandvik PO, Zhao P, Zhang L, Shen J, Bala MM, Sohani ZN, Wong E, Busse JW, Ebrahim S, Malaga G, Rios LP, Wang Y, Chen Q, Guyatt GH, Sun X. Dipeptidyl peptidase-4 inhibitors and risk of heart failure in type 2 diabetes: systematic review and meta-analysis of randomised and observational studies. *BMJ*. 2016;352:i610.
- Brandt I, Lambeir AM, Ketelslegers JM, Vanderheyden M, Scharpe S, De Meester I. Dipeptidyl-peptidase IV converts intact B-type natriuretic peptide into its des-SerPro form. *Clin Chem*. 2006;52(1):82–7.
- Costello-Boerrigter LC, Boerrigter G, Redfield MM, Rodeheffer RJ, Urban LH, Mahoney DW, Jacobsen SJ, Heublein DM, Burnett JC Jr. Amino-terminal pro-B-type natriuretic peptide and B-type natriuretic peptide in the general community: determinants and detection of left ventricular dysfunction. *J Am Coll Cardiol*. 2006;47(2):345–53.
- Yoshimura M, Yasue H, Morita E, Sakaino N, Jougasaki M, Kurose M, Mukoyama M, Saito Y, Nakao K, Imura H. Hemodynamic, renal, and hormonal responses to brain natriuretic peptide infusion in patients with congestive heart failure. *Circulation*. 1991;84(4):1581–8.
- Plante E, Menaouar A, Danalache BA, Broderick TL, Jankowski M, Gutkowska J. Treatment with brain natriuretic peptide prevents the development of cardiac dysfunction in obese diabetic db/db mice. *Diabetologia*. 2014;57(6):1257–67.
- Gruden G, Landi A, Bruno G. Natriuretic peptides, heart, and adipose tissue: new findings and future developments for diabetes research. *Diabetes Care*. 2014;37(11):2899–908.
- Takahashi A, Ihara M, Yamazaki S, Asanuma H, Asakura M, Kitakaze M. Impact of either GLP-1 agonists or DPP-4 inhibitors on pathophysiology of heart failure. *Int Heart J*. 2015;56(4):372–6.
- Mair J. Biochemistry of B-type natriuretic peptide—where are we now? *Clin Chem Lab Med*. 2008;46(11):1507–14.

13. Alonso N, Lupon J, Barallat J, de Antonio M, Domingo M, Zamora E, Moliner P, Galan A, Santesmases J, Pastor C, Mauricio D, Bayes-Genis A. Impact of diabetes on the predictive value of heart failure biomarkers. *Cardiovasc Diabetol*. 2016;15(1):151.
14. von Scholten BJ, Reinhard H, Hansen TW, Lindhardt M, Petersen CL, Wiinberg N, Hansen PR, Parving HH, Jacobsen PK, Rossing P. Additive prognostic value of plasma N-terminal pro-brain natriuretic peptide and coronary artery calcification for cardiovascular events and mortality in asymptomatic patients with type 2 diabetes. *Cardiovasc Diabetol*. 2015;14:59.
15. Niederkofler EE, Kiernan UA, O'Rear J, Menon S, Saghir S, Protter AA, Nelson RW, Schellenberger U. Detection of endogenous B-type natriuretic peptide at very low concentrations in patients with heart failure. *Circ Heart Fail*. 2008;1(4):258–64.
16. Seferian KR, Tamm NN, Semenov AG, Mukharyamova KS, Tolstaya AA, Koshkina EV, Kara AN, Krasnoselsky MI, Apple FS, Esakova TV, Filatov VL, Katrukha AG. The brain natriuretic peptide (BNP) precursor is the major immunoreactive form of BNP in patients with heart failure. *Clin Chem*. 2007;53(5):866–73.
17. Heublein DM, Huntley BK, Boerrigter G, Cataliotti A, Sandberg SM, Redfield MM, Burnett JC Jr. Immunoreactivity and guanosine 3',5'-cyclic monophosphate activating actions of various molecular forms of human B-type natriuretic peptide. *Hypertension*. 2007;49(5):1114–9.
18. Boerrigter G, Costello-Boerrigter LC, Harty GJ, Lapp H, Burnett JC Jr. Deserine-proline brain natriuretic peptide 3-32 in cardiorenal regulation. *Am J Physiol Regul Integr Comp Physiol*. 2007;292(2):R897–901.
19. Scirica BM, Braunwald E, Raz I, Cavender MA, Morrow DA, Jarolim P, Udell JA, Mosenzon O, Im K, Umez-Eronini AA, Pollack PS, Hirshberg B, Frederich R, Lewis BS, McGuire DK, Davidson J, Steg PG, Bhatt DL. Heart failure, saxagliptin, and diabetes mellitus: observations from the SAVOR-TIMI 53 randomized trial. *Circulation*. 2015;132(15):e198.
20. Zannad F, Cannon CP, Cushman WC, Bakris GL, Menon V, Perez AT, Fleck PR, Mehta CR, Kupfer S, Wilson C, Lam H, White WB. Heart failure and mortality outcomes in patients with type 2 diabetes taking alogliptin versus placebo in EXAMINE: a multicentre, randomised, double-blind trial. *Lancet*. 2015;385(9982):2067–76.
21. Fadini GP, Bonora BM, Cappellari R, Menegazzo L, Vedovato M, Iori E, Marescotti MC, Albiero M, Avogaro A. Acute effects of linagliptin on progenitor cells, monocyte phenotypes, and soluble mediators in type 2 diabetes. *J Clin Endocrinol Metab*. 2016;101(2):748–56.
22. von Websky K, Reichetzeder C, Hocher B. Physiology and pathophysiology of incretins in the kidney. *Curr Opin Nephrol Hypertens*. 2014;23(1):54–60.
23. Hocher B, Reichetzeder C, Alter ML. Renal and cardiac effects of DPP4 inhibitors—from preclinical development to clinical research. *Kidney Blood Press Res*. 2012;36(1):65–84.
24. McAlister FA, Ezekowitz J, Tonelli M, Armstrong PW. Renal insufficiency and heart failure: prognostic and therapeutic implications from a prospective cohort study. *Circulation*. 2004;109(8):1004–9.
25. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, Kusek JW, Eggers P, Van Lente F, Greene T, Coresh J. A new equation to estimate glomerular filtration rate. *Ann Intern Med*. 2009;150(9):604–12.
26. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis*. 2002;39(2 Suppl 1):S1–266.
27. Poncina N, Albiero M, Menegazzo L, Cappellari R, Avogaro A, Fadini GP. The dipeptidyl peptidase-4 inhibitor saxagliptin improves function of circulating pro-angiogenic cells from type 2 diabetic patients. *Cardiovasc Diabetol*. 2014;13:92.
28. Kim HN, Januzzi JL Jr. Natriuretic peptide testing in heart failure. *Circulation*. 2011;123(18):2015–9.
29. Scirica BM, Bhatt DL, Braunwald E, Steg PG, Davidson J, Hirshberg B, Ohman P, Frederich R, Wiviott SD, Hoffman EB, Cavender MA, Udell JA, Desai NR, Mosenzon O, McGuire DK, Ray KK, Leiter LA, Raz I. Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med*. 2013;369(14):1317–26.
30. Fadini GP, Avogaro A, Degli Esposti L, Russo P, Saragoni S, Buda S, Rosano G, Pecorelli S, Pani L. Risk of hospitalization for heart failure in patients with type 2 diabetes newly treated with DPP-4 inhibitors or other oral glucose-lowering medications: a retrospective registry study on 127,555 patients from the Nationwide OsMed Health-DB Database. *Eur Heart J*. 2015;36(36):2454–62.
31. Chang CH, Chang YC, Lin JW, Caffrey JL, Wu LC, Lai MS, Chuang LM. No increased risk of hospitalization for heart failure for patients treated with dipeptidyl peptidase-4 inhibitors in Taiwan. *Int J Cardiol*. 2016;220:14–20.
32. Ou S-M, Chen H-T, Kuo S-C, Chen T-J, Shih C-J, Chen Y-T. Dipeptidyl peptidase-4 inhibitors and cardiovascular risks in patients with pre-existing heart failure. *Heart*. 2016. doi:10.1136/heartjnl-2016-309687.
33. Sato A, Yoshihisa A, Kanno Y, Takiguchi M, Miura S, Shimizu T, Nakamura Y, Yamauchi H, Owada T, Sato T, Suzuki S, Oikawa M, Yamaki T, Sugimoto K, Kunii H, Nakazato K, Suzuki H, Saitoh SI, Takeishi Y. Associations of dipeptidyl peptidase-4 inhibitors with mortality in hospitalized heart failure patients with diabetes mellitus. *ESC Heart Fail*. 2016;3(2):77–85.
34. White WB, Cannon CP, Heller SR, Nissen SE, Bergenstal RM, Bakris GL, Perez AT, Fleck PR, Mehta CR, Kupfer S, Wilson C, Cushman WC, Zannad F. Alogliptin after acute coronary syndrome in patients with type 2 diabetes. *N Engl J Med*. 2013;369(14):1327–35.
35. Chaykovska L, von Websky K, Rahnenfuhrer J, Alter M, Heiden S, Fuchs H, Runge F, Klein T, Hocher B. Effects of DPP-4 inhibitors on the heart in a rat model of uremic cardiomyopathy. *PLoS ONE*. 2011;6(11):e27861.
36. Hocher B, Sharkovska Y, Mark M, Klein T, Pfab T. The novel DPP-4 inhibitors linagliptin and BI 14361 reduce infarct size after myocardial ischemia/reperfusion in rats. *Int J Cardiol*. 2013;167(1):87–93.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

