

Editorial

Open Access

Which is the best lipid-modifying strategy in metabolic syndrome and diabetes: fibrates, statins or both?

Alexander Tenenbaum^{1,2} and Enrique Z Fisman*^{1,2}

Address: ¹Cardiac Rehabilitation Institute, the Chaim Sheba Medical Center, 52621 Tel-Hashomer, Ramat-Gan, Israel and ²Sackler Faculty of Medicine, Tel-Aviv University, 69978 Ramat-Aviv, Tel-Aviv, Israel

Email: Alexander Tenenbaum - altenen@post.tau.ac.il; Enrique Z Fisman* - zfisman@post.tau.ac.il

* Corresponding author

Published: 01 December 2004

Received: 01 December 2004

Cardiovascular Diabetology 2004, **3**:10 doi:10.1186/1475-2840-3-10

Accepted: 01 December 2004

This article is available from: <http://www.cardiab.com/content/3/1/10>

© 2004 Tenenbaum and Fisman; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Although less clinical intervention studies have been performed with fibrates than with statins, there are evidences indicating that fibrates may reduce risk of cardiovascular events. The potential clinical benefit of the fenofibrate will be specified by the ongoing Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study, which rationale, methods and aims have been just published.

Controlled clinical trials show similar or even greater cardiovascular benefits from statins-based therapy in patient subgroups with diabetes compared with overall study populations. Therefore, statins are the drug of first choice for aggressive lipid lowering actions and reducing risk of coronary artery disease in these patients. However, current therapeutic use of statins as monotherapy is still leaving many patients with mixed atherogenic dyslipidemia at high risk for coronary events. A combination statin/fibrate therapy may be often necessary to control all lipid abnormalities in patients with metabolic syndrome and diabetes adequately, since fibrates provide additional important benefits, particularly on triglyceride and HDL-cholesterol levels. Thus, this combined therapy concentrates on all the components of the mixed dyslipidemia that often occurs in persons with diabetes or metabolic syndrome, and may be expected to reduce cardiovascular morbidity and mortality.

Safety concerns about some fibrates such as gemfibrozil may lead to exaggerate precautions regarding fibrate administration and therefore diminish the use of the seagents. However, other fibrates, such as bezafibrate and fenofibrate appear to be safer and better tolerated. We believe that a proper co-administration of statins and fibrates, selected on basis of their safety, could be more effective in achieving a comprehensive lipid control as compared with monotherapy.

Due to their beneficial effects on glucose and lipid metabolism, peroxisome proliferator activated receptors (PPAR's) alpha agonists (fibrates) are good potential candidates for reducing the risk of myocardial infarction (MI)

in subjects with metabolic syndrome [1-3]. Although less clinical intervention studies have been performed with fibrates than with statins, there are evidences indicating that fibrates may reduce risk of cardiovascular disease and

particularly non-fatal MI [4-10]. Interestingly, reduction of cardiovascular disease with one of the fibric acid derivatives – gemfibrozil – was more pronounced in patients displaying baseline characteristics very similar to metabolic syndrome definitions [4,5].

There have been no direct head-to-head comparisons of a statin with a fibrate in any clinical endpoint trial. However, compared with statins, fibrates appear to more selectively target the therapeutic goals in obese individuals with features of insulin resistance and metabolic syndrome (i.e. with near-goal low-density lipoprotein (LDL)-cholesterol and inappropriate high-density lipoprotein (HDL)-cholesterol and triglyceride levels).

The primary-prevention trial Helsinki Heart Study showed that treatment with gemfibrozil led to a significant reduction in major cardiovascular events [4]. Regarding secondary prevention, in the VAHIT study (Veterans Affairs High-density lipoprotein cholesterol Intervention Trial) – which included 30% of diabetic patients – gemfibrozil reduced the occurrence of major cardiovascular events by 22 % [5]. Similarly, reduction of cardiovascular disease with gemfibrozil was more pronounced in patients displaying above three of the features of metabolic syndrome [11,12].

In two previous small studies bezafibrate decreased the rate of progression of coronary atherosclerosis and decreased coronary event rate [6,7]. In another large trial in 1568 men with lower extremity arterial disease with a relatively short follow-up period, bezafibrate reduced the severity of intermittent claudication for up to three years. [8]. In general, the incidence of coronary heart disease in patients on bezafibrate has tended to be lower, but this tendency did not reach statistical significance. However, bezafibrate had significantly reduced the incidence of non-fatal coronary events, particularly in those aged <65 years at entry, in whom all coronary events may also be reduced [8]. In the Bezafibrate Infarction Prevention (BIP) study an overall trend of a 9.4% reduction of the incidence of primary end point (fatal or non-fatal myocardial infarction or sudden death) was observed. The reduction in the primary end point in 459 patients with high baseline triglycerides (≥ 200 mg/dL) was significant [9]. These results are consistent with studies in experimental models showing that pre-treatment of rats with the PPAR-alpha agonist clofibrate causes a significant reduction in induced myocardial infarct size of 43% [13].

Recently, reduced incidence of type 2 diabetes in patients with impaired fasting glucose level on bezafibrate has been demonstrated [14]. The potential clinical benefit of the other widespread fibric acid derivative, fenofibrate, on the reduction of cardiovascular disease is still unknown

and will be specified by the ongoing Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study, which rationale, methods and aims have been just published [15]. It will be the largest (approximately 10000 patients) ever conducted fibrate-based controlled clinical trial in diabetic patients. The results are expected for 2005. An added strength of this trial is its ability to examine important clinical outcomes across diverse ethnic and gender subgroups. The results of this study will clarify whether the beneficial lipid-modifying effects of micronised fenofibrate lead to a reduction of cardiovascular morbidity and mortality.

Despite increasing use of statins, a significant number of coronary events still occur and many of such events take place in patients presenting with the metabolic syndrome. Whereas statins remain the drug of choice for patients who need to achieve the LDL-cholesterol goal, fibrate therapy may represent the alternative intervention for subjects with atherogenic dyslipidemia typical for metabolic syndrome and an LDL-cholesterol already close to goal values. In addition, the concomitant use of fibrates seems to be attractive in patients whose LDL-cholesterol is controlled by statin therapy but whose HDL-cholesterol and/or triglyceride levels are still inappropriate [16-19]. This strategy will be tested in the ongoing Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial [20].

The factor that dominates in overweight-related metabolic syndrome is the permanent elevation of plasma free fatty acids (FFA) and the predominant utilization of lipids by the muscle inducing a diminution of glucose uptake and insulin resistance. Currently, an insulin-resistant state – as the key phase of metabolic syndrome – constitutes the major risk factor for development of macrovascular complications [21-23]. On the basis of the current concept of the evolution of adipogenesis via PPAR modulation toward insulin resistance and atherothrombotic macrovascular complications (including MI), the decreasing of plasma FFA and improving of insulin sensitization by PPAR agonists seems to be a logical and valuable goal for therapy.

It is important to note that on a whole-body level, lipid and glucose metabolisms interact intimately. Briefly, PPAR alpha is activated by fibric acids (e.g. bezafibrate) and form heterodimers with the 9-cis retinoic acid receptor (RXR). These heterodimers bind to peroxisome proliferator response elements, which are located in numerous gene promoters and increase the level of the expression of mRNAs encoded by PPAR alpha target genes. Bezafibrate reduces triglyceride plasma levels through increases in the expression of genes involved in fatty acid-beta oxidation and by decrease in apolipoprotein C-III gene expression. Fibric acids increase HDL-cholesterol partly by increasing

apolipoprotein A-I and apolipoprotein A-II gene expression. Their triglyceride-lowering and HDL-cholesterol raising effects lead to decreased systemic availability of fatty acid, diminished fatty acid uptake by muscle with improvement of insulin sensitization and reduced plasma glucose level [24-28].

Evidence also suggests that there is a 'fibrate effect' that mediates the reduction in CHD risk beyond the favorable impact of these agents on HDL-cholesterol levels. This last notion is consistent with the pleiotropic effects of fibrates which are known to be related to their mechanisms of action [29]. Being PPAR alpha ligands, fibrates have a significant impact on the synthesis of several apolipoproteins (apo) and enzymes of lipoprotein metabolism as well as on the expression of several genes involved in fibrinolysis and inflammation. Such changes contribute to improve the catabolism of triglyceride-rich lipoproteins, leading to a substantial increase in HDL-cholesterol levels accompanied by a shift in the size and density of LDL particles (from small, dense LDL particles to larger, more buoyant cholesteryl ester-rich LDL).

Controlled clinical trials show similar or even greater cardiovascular benefits from statins-based therapy in patient subgroups with diabetes, impaired fasting glucose, and metabolic syndrome, compared with overall study populations. Therefore, statins are the drug of first choice for aggressive lipid lowering actions and reducing risk of coronary artery disease in these patients. However, current therapeutic use of statins as monotherapy is still leaving many patients with mixed atherogenic dyslipidemia at high risk for coronary events.

A combination statin/fibrate therapy may be often necessary to control all lipid abnormalities in patients with metabolic syndrome and diabetes adequately, since fibrates provide additional important benefits, particularly on triglyceride and HDL-C levels. Thus, this combined therapy concentrates on all the components of the mixed dyslipidemia that often occurs in persons with diabetes or metabolic syndrome, and may be expected to reduce cardiovascular morbidity and mortality.

Safety concerns about some fibrates such as gemfibrozil may lead to exaggerate precautions regarding fibrate administration and therefore diminish the use of the agents. However, other fibrates such as bezafibrate and fenofibrate appear to be safer and better tolerated [30-36]. We believe that a proper co-administration of statins and fibrates, selected on basis of their safety, could be more effective in achieving a comprehensive lipid control as compared with monotherapy.

Competing interests

The author(s) declare that they have no competing interests.

Acknowledgements

This work was supported in part by the Cardiovascular Diabetology Research Foundation (RA 58-040-684-1), Holon, Israel, and the Research Authority of Tel-Aviv University (Citernick grant 01250239).

References

1. Sacks FM, for the Expert Group on HDL Cholesterol: **The role of high-density lipoprotein [HDL] cholesterol in the prevention and treatment of coronary heart disease: Expert Group recommendations.** *Am J Cardiol* 2002, **90**:139-143.
2. Fruchart JC: **Peroxisome proliferator-activated receptor-alpha activation and high-density lipoprotein metabolism.** *Am J Cardiol* 2001, **88(12A)**:24N-29N.
3. Verges B: **Clinical interest of PPARs ligands.** *Diabetes Metab* 2004, **30**:7-12.
4. Frick MH, Elo O, Haapa K, Heinonen OP, Heinsalmi P, Helttunen JK, Kaitaniemi P, Koskinen P, Manninen V: **Helsinki Heart Study: primary-prevention trial with gemfibrozil in middle-aged men with dyslipidemia. Safety of treatment, changes in risk factors, and incidence of coronary heart disease.** *N Engl J Med* 1987, **317**:1237-1245.
5. Rubins HB, Robins SJ, Collins D, Fye CL, Anderson JW, Elam MB, Faas FH, Linares E, Schaefer EJ, Schectman G, Wilt TJ, Wittes J: **Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. Veterans Affairs High-Density Lipoprotein Cholesterol Intervention Trial Study Group.** *N Engl J Med* 1999, **341**:410-418.
6. Ericsson CG, Nilsson J, Grip L, Svane B, Hamsten A: **Effect of bezafibrate treatment over five years on coronary plaques causing 20% to 50% diameter narrowing (The Bezafibrate Coronary Atherosclerosis Intervention Trial (BECAIT)).** *Am J Cardiol* 1997, **80**:1125-1129.
7. Elkees RS, Diamond JR, Poulter C, Dhanjil S, Nicolaidis AN, Mahmood S, Richmond W, Mather H, Sharp P, Feher MD: **Cardiovascular outcomes in type 2 diabetes. A double-blind placebo-controlled study of bezafibrate: the St. Mary's, Ealing, Northwick Park Diabetes Cardiovascular Disease Prevention (SENDCAP) Study.** *Diabetes Care* 1998, **21**:641-648.
8. Meade T, Zuhrie R, Cook C, Cooper J: **Bezafibrate in men with lower extremity arterial disease: randomised controlled trial.** *BMJ* 2002, **325(7373)**:1139.
9. **Secondary prevention by raising HDL cholesterol and reducing triglycerides in patients with coronary artery disease: the Bezafibrate Infarction Prevention (BIP) study.** *Circulation* 2000, **102**:21-27.
10. Diabetes Atherosclerosis Intervention Study Investigators: **Effect of fenofibrate on progression of coronary-artery disease in type 2 diabetes: the diabetes atherosclerosis intervention study, a randomised study.** *Lancet* 2001, **357**:905-910.
11. Manninen V, Tenkanen L, Koskinen P, Huttunen JK, Manttari M, Heinonen OP, Frick MH: **Joint effects of serum triglyceride and LDL cholesterol and HDL cholesterol concentrations on coronary heart disease risk in the Helsinki Heart Study. Implications for treatment.** *Circulation* 1992, **85**:37-45.
12. Rubins HB, Robins SJ, Collins D, Nelson DB, Elam MB, Schaefer EJ, Faas FH, Anderson JW: **Diabetes, plasma insulin, and cardiovascular disease: subgroup analysis from the Department of Veterans Affairs high-density lipoprotein intervention trial (VA-HIT).** *Arch Intern Med* 2002, **162**:2597-2604.
13. Wayman NS, Ellis BL, Thiemermann C: **Ligands of the peroxisome proliferator-activated receptor-PPAR-a reduce myocardial infarct size.** *Med Sci Monit* 2002, **8(7)**:BR243-247.
14. Tenenbaum A, Motro M, Fisman EZ, Schwammenthal E, Adler Y, Goldenberg I, Leor J, Boyko V, Mandelzweig L, Behar S: **Peroxisome proliferator-activated receptors ligand bezafibrate for prevention of type 2 diabetes mellitus in patients with coronary artery disease.** *Circulation* 2004, **109**:2197-2202.

15. Keech AC, the FIELD Study Investigators: **The need for a large-scale trial of fibrate therapy in diabetes: the rationale and design of the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study.** ISRCTN64783481. *Cardiovasc Diabetol* 2004, **3**:9.
16. Robins SJ: **Cardiovascular disease with diabetes or the metabolic syndrome: should statins or fibrates be first line lipid therapy?** *Curr Opin Lipidol* 2003, **14**:575-583.
17. Fazio S, Linton MF: **The role of fibrates in managing hyperlipidemia: mechanisms of action and clinical efficacy.** *Curr Atheroscler Rep* 2004, **6**:148-157.
18. **Role of fibrates in reducing coronary risk: a UK Consensus.** *Curr Med Res Opin* 2004, **20**:241-247.
19. Tenenbaum A, Fisman EZ, Motro M: **Bezafibrate and simvastatin: different beneficial effects for different therapeutic aims.** *J Clin Endocrinol Metab* 2004, **89**:1978.
20. Prisant LM: **Clinical trials and lipid guidelines for type II diabetes.** *J Clin Pharmacol* 2004, **44**:423-430.
21. Groop LC: **Insulin resistance: the fundamental trigger of type 2 diabetes.** *Diabetes Obes Metab* 1999;S1-S7.
22. Tenenbaum A, Fisman EZ, Motro M: **Metabolic syndrome and type 2 diabetes mellitus: focus on peroxisome proliferator activated receptors (PPAR).** *Cardiovasc Diabetol* 2003, **2**:4.
23. Lakka HM, Laaksonen DE, Lakka TA, Niskanen LK, Kumpusalo E, Tuomilehto J, Salonen JT: **The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men.** *JAMA* 2002, **288**:2709-2716.
24. Fruchart JC, Staels B, Duriez P: **The role of fibric acids in atherosclerosis.** *Curr Atheroscler Rep* 2001, **3**:83-92.
25. Rovellini A, Sommariva D, Branchi A, Maraffi F, Montalto C, Gandini R, Fasoli A: **Effects of slow release bezafibrate on the lipid pattern and on blood glucose of type 2 diabetic patients with hyperlipidaemia.** *Pharmacol Res* 1992, **25**:237-245.
26. Taniguchi A, Fukushima M, Sakai M, Tokuyama K, Nagata I, Fukunaga A, Kishimoto H, Doi K, Yamashita Y, Matsuura T, Kitatani N, Okumura T, Nagasaka S, Nakaishi S, Nakai Y: **Effects of bezafibrate on insulin sensitivity and insulin secretion in non-obese Japanese type 2 diabetic patients.** *Metabolism* 2001, **50**:477-480.
27. Jonkers IJ, Mohrschlatt MF, Westendorp RG, van der Laarse A, Smelt AH: **Severe hypertriglyceridemia with insulin resistance is associated with systemic inflammation: reversal with bezafibrate therapy in a randomized controlled trial.** *Am J Med* 2002, **112**:275-280.
28. Jones IR, Swai A, Taylor R, Miller M, Laker MF, Alberti KG: **Lowering of plasma glucose concentrations with bezafibrate in patients with moderately controlled NIDDM.** *Diabetes Care* 1990, **13**:855-863.
29. Despres JP, Lemieux I, Robins SJ: **Role of fibric Acid derivatives in the management of risk factors for coronary heart disease.** *Drugs* 2004, **64**(19):2177-2198.
30. Jokubaitis LA: **Fluvastatin in combination with other lipid-lowering agents.** *Br J Clin Pract Suppl* 1996, **77A**:28-32.
31. Gavish D, Leibovitz E, Shapira I, Rubinstein A: **Bezafibrate and simvastatin combination therapy for diabetic dyslipidaemia: efficacy and safety.** *J Intern Med* 2000, **247**:563-569.
32. Kyrklund C, Backman JT, Kivisto KT, Neuvonen M, Laitila J, Neuvonen PJ: **Plasma concentrations of active lovastatin acid are markedly increased by gemfibrozil but not by bezafibrate.** *Clin Pharmacol Ther* 2001, **69**:340-345.
33. Beggs PW, Clark DW, Williams SM, Coulter DM: **A comparison of the use, effectiveness and safety of bezafibrate, gemfibrozil and simvastatin in normal clinical practice using the New Zealand Intensive Medicines Monitoring Programme (IMMP).** *Br J Clin Pharmacol* 1999, **47**:99-104.
34. Farnier M, Salko T, Isaacsohn JL, Troendle AJ, Dejager S, Gonasun L: **Effects of baseline level of triglycerides on changes in lipid levels from combined fluvastatin + fibrate (bezafibrate, fenofibrate, or gemfibrozil).** *Am J Cardiol* 2003, **92**:794-797.
35. Farnier M: **Combination therapy with an HMG-CoA reductase inhibitor and a fibric acid derivative: a critical review of potential benefits and drawbacks.** *Am J Cardiovasc Drugs* 2003, **3**:169-178.
36. Shek A, Ferrill MJ: **Statin-fibrate combination therapy.** *Ann Pharmacother* 2001, **35**:908-917.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

