

CORRECTION

Open Access



Correction to: Type 2 diabetes-associated carotid plaque burden is increased in patients with retinopathy compared to those without retinopathy

Núria Alonso¹, Alicia Traveset², Esther Rubinat^{3,4}, Emilio Ortega⁵, Nuria Alcubierre⁴, Jordi Sanahuja⁷, Marta Hernández^{3,6}, Angels Betriu⁶, Carmen Jurjo², Elvira Fernández^{6,8} and Didac Mauricio^{1*}

Correction to: *Cardiovasc Diabetol* (2015) 14:33

<https://doi.org/10.1186/s12933-015-0196-1>

The author found errors in Table 1 after publication of the original article [1].

The corrected Table 1 is presented in this erratum.

Table 1 Characteristics of the study groups

	Diabetes without retinopathy (n = 159)	Diabetic with retinopathy (n = 153)	p
Sex—men, n (%)	82 (51.6%)	76 (49.7%)	0.824
Non caucasian, n (%)	7 (4.40%)	7 (4.58%)	1.000
Age (years)	59.0 [48.5;66.0]	61.0 [54.0;68.0]	0.012
Disease duration (years)	6.00 [2.50;10.0]	11.0 [6.00;20.0]	< 0.001
Insulin treatment, n (%)	20 (12.6%)	84 (54.9%)	< 0.001
Smoking (current/past/never) (n)	32/55/71	30/44/79	0.442
Antiplatelet treatment, n (%)	47 (29.6%)	71 (46.4%)	0.003
Dyslipidaemia, n (%)	67 (42.1%)	78 (51.0%)	0.147
Hypertension, n (%)	80 (50.3%)	103 (67.3%)	0.003
Systolic BP (mmHg)	134 [123;145]	143 [133;159]	< 0.001
Diastolic BP (mmHg)	76.0 [70.5;83.0]	77.5 [68.8;85.2]	0.752
Heart rate (b/min)	75.0 [68.0;82.0]	77.0 [70.0;86.0]	0.076
Body mass index (kg/m ²)	30.3 [27.4;34.0]	31.1 [28.3;35.0]	0.247
Waist (cm)	103 [96.8;111]	107 [101;114]	0.007
Haemoglobin A1c (%)	7.00 [6.50;7.90]	8.10 [7.20;9.10]	< 0.001
Total-cholesterol (mg/dL)	184 [163;206]	181 [163;204]	0.988
HDL-cholesterol (mg/dL)	47.0 [40.0;57.0]	50.5 [42.0;60.2]	0.03
LDL-cholesterol (mg/dL)	109 [90.4;130]	105 [86.5;128]	0.268
Triglycerides (mg/dL)	118 [89.5;169]	116 [83.0;168]	0.617
Creatinine (mg/dL)	0.79 [0.69;0.92]	0.79 [0.68;0.93]	0.797
Urinary albumin/creatinine ratio (mg/g)	5.80 [3.19;10.9]	12.4 [6.00;32.7]	< 0.001

Data are presented as the median values and interquartile ranges

BP blood pressure

*Correspondence: didacmauricio@gmail.com

¹ Department of Endocrinology and Nutrition, Health Sciences Research Institute and University Hospital Germans Trias i Pujol, Badalona, Spain
Full list of author information is available at the end of the article



© The Author(s) 2018. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Author details

¹ Department of Endocrinology and Nutrition, Health Sciences Research Institute and University Hospital Germans Trias i Pujol, Badalona, Spain. ² Department of Ophthalmology, Hospital Universitari Arnau de Vilanova, Lleida, Spain. ³ Department of Endocrinology and Nutrition, Hospital Universitari Arnau de Vilanova, Lleida, Spain. ⁴ Institut de Recerca Biomedica de Lleida, University of Lleida, Lleida, Spain. ⁵ Department of Endocrinology and Nutrition, Institut d'Investigacions Biomediques August Pi Suñer, CIBER de Diabetes y Enfermedades Metabólicas asociadas, Hospital Clinic, 08036 Barcelona, Spain. ⁶ Unitat de Detecció i Tractament de Malalties Aterotrombòtiques, Hospital Universitari Arnau de Vilanova, Lleida, Spain. ⁷ Department of Neurology, Hospital Universitari Arnau de Vilanova, Lleida, Spain. ⁸ Department of Nephrology, Hospital Universitari Arnau de Vilanova, Lleida, Spain.

The original article can be found online at <https://doi.org/10.1186/s12933-015-0196-1>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 29 March 2018 Accepted: 29 March 2018

Published online: 05 April 2018

Reference

1. Alonso N, Traveset A, Rubinat E, Ortega E, Alcubierre N, Sanahuja J, Hernández M, Betriu A, Jurjo C, Fernández E, Mauricio D. Type 2 diabetes-associated carotid plaque burden is increased in patients with retinopathy compared to those without retinopathy. *Cardiovasc Diabetol*. 2015;14:33. <https://doi.org/10.1186/s12933-015-0196-1>.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

